

Contribution to the knowledge of the world Rhynchitidae (Coleoptera)

Andrei A. Legalov

Legalov A.A. 2009. Contribution to the knowledge of the world Rhynchitidae (Coleoptera). *Baltic J. Coleopterol.* 9(1): 55 - 88.

New genus *Pilosauletes* Legalov, gen.n (type species: *Auletobius aurichalceus* Voss, 1939) and new species *Pseudominurus bananensis* Legalov, sp.n. (Zaire), *Pseudomesauletes kryzhanovskyi* Legalov, sp.n. (Fujian, Yunnan), *P. poirasi* Legalov, sp.n. (India, Nepal), *Heterorhynchites macros* Legalov, sp.n. (Sabah) and *H. perakensis* Legalov, sp.n. (Malaysia: Perak) are described. New synonyms: *Deporaus paucus* Voss, 1941, **syn.n.** for *Caenorhinus minimus koreanus* (Voss, 1929); *Rhynchites bicuspis* Voss, 1924, **syn.n.** for *Auletomorphus cupidio* (Pascoe, 1875); *Rhynchites rhodesianus* Voss, 1938, **syn.n.**, *Rhynchites natalensis* Voss, 1938, **syn.n.**, *Afrovolvulus katonensis* Legalov, 2004, **syn.n.** for *Afrorhynchites bipubescens* (Hustache, 1929); *Callirhynchites biumbanus* Legalov, 2007, **syn.n.** for *Callirhynchites mundus* (Voss, 1938); *Rhynchites kirishenkoi* Ter-Minassian, 1944, **syn.n.** for *Thompsonirhinus mandschuricus* (Voss, 1939); *Byctiscus chinensis* Formanek, 1911, **syn.n.** for *B. impressus* (Fairmaire, 1899). New systematic placements: *Auletobius pallidus* Voss, 1933 from subgenus *Auletobius* s. str. to subgenus *Auletobioides* Legalov, 2007; *Auletobius aurichalceus* Voss, 1939, **placem.n.** from subtribe *Auletobiina* Legalov, 2001 to subtribe *Mandelschtamiina* Legalov, 2003; *Gymnauletes* Legalov, 2001, **placem.n.** (type species: *Auletes glaber* Faust, 1892) and *Gymnauletobius* Legalov, 2007, **placem.n.** (type species: *Auletobius nitidus* Voss, 1922) from subtribe *Pseudauletinina* Voss, 1933 to subtribe *Pseudomesauletina* Legalov, 2003; *Pseudomesauletes contristatus* (Voss, 1939), **placem.n.** and *P. ueleanus* (Voss, 1939), **placem.n.** from genus *Alonsoiauletes* Legalov, 2003 to genus *Pseudomesauletes* Legalov, 2001; *Pseudomesauletes viridimicans* (Voss, 1939), **placem.n.** from subgenus *Metallauletes* Legalov, 2007 to subgenus *Afromesauletes* Legalov, 2003; *Cateugnamptus hirsutus* (Voss, 1924), **placem.n.** from genus *Neoeugnamptus* Legalov, 2003 to genus *Cateugnamptus* Legalov, 2007; *Eusproda tumida* (Voss, 1938), **placem.n.** from genus *Exrhynchites* Voss, 1930 to genus *Eusproda* Sawada, 1987; *Caenorhinus gilviventris* (Voss, 1938), **placem.n.** from *Caenorhinus incertae sedis* to subgenus *Flavodeporaus* Legalov, 2007; *Proelautobius erythropterus* (Voss, 1921), **placem.n.** from genus *Rubrinvolvulus* Legalov, 2003 to genus *Proelautobius* Legalov, 2007; *Afrorhynchites conradti* (Voss, 1938), **placem.n.** from genus *Pararhynchites* Legalov, 2003 to genus *Afrorhynchites* Legalov, 2003; *Callirhynchites mundus* (Voss, 1938), **placem.n.** from genus *Pararhynchites* Legalov, 2003 to genus *Callirhynchites* Voss, 1938; *Thompsonirhinus mandschuricus* (Voss, 1939), **placem.n.** from genus *Parinvolvulus* Legalov, 2003 to genus *Thompsonirhinus* Legalov, 2003. *Pseudominurus mubendensis* Legalov, 2007 in fauna of Zaire and *Yunnanuletes heishuensis* Legalov, 2007 in fauna of Sichuan were proven for the first time. Nine new combinations are established. Type specimens of the following

species were studied by the author: *Auletobius diversicolor* Voss, 1939, *Auletobius kraatzi* Voss, 1922, *Auletobius urundiensis* Voss, 1939, *Auletobius aurichalceus* Voss, 1939, *Auletobius brevihiatus* Voss, 1939, *Auletobius pallidus* Voss, 1933, *Pseudauleutes nitens* Voss, 1930, *Auletobius bicolor* Voss, 1922, *Auletobius castaneus* Voss, 1930, *Auletobius cognatus* Voss, 1930, *Auletobius glaber* Faust, 1892, *Auletobius contristatus* Voss, 1939, *Auletobius ueleanus* Voss, 1939, *Auletobius consimilis* Voss, 1930, *Auletobius kuntzeni* Voss, 1922, *Auletobius togoensis* f. *viridimicans* Voss, 1939, *Eugnamptus diversus* Voss, 1948, *Eugnamptus inclusus* Voss, 1941, *Anthribus collaris* Fabricius, 1801, *Attelabus tristis* Fabricius, 1794, *Deporaus scolocnemoides* Voss, 1935, *Deporaus tumidus* Voss, 1938, *Deporaus pauculus* Voss, 1941, *Deporaus kolbei* Voss, 1938, *Deporaus gilviventris* Voss, 1938, *Attelabus planirostris* Fabricius, 1801, *Curculio bicolor* Fabricius, 1775, *Rhynchites bisulcatus* Voss, 1921, *Coenorrhinus decumanus* Voss, 1930, *Rhynchites bicuspidis* Voss, 1924, *Rhynchites similatus* Voss, 1938, *Rhynchites erythropterus* Voss, 1921, *Rhynchites castaneus* Jekel, 1860, *Rhynchites conradti* Voss, 1938, *Rhynchites natalensis* Voss, 1938, *Rhynchites mundus* Voss, 1938, *Rhynchites lenaeus* ssp. *slovenicus* Purkyne, 1954, *Rhynchites indubius* Voss, 1930, *Rhynchites mandschuricus* Voss, 1939, *Rhynchites balaninoides* Voss, 1938, *Attelabus betuleti* Fabricius, 1792, *Byctiscus chinensis* Formanek, 1911, *Byctiscus impressus* f. *thibetana* Voss, 1933.

Key words: Coleoptera, Curculionoidea, Rhynchitidae, Attelabidae, new genus, new species, new placement, new combination

Andrei Legalov. Zoological Museum, Institute of Animal Systematics and Ecology, SB RAS, Frunze street-11, Novosibirsk 630091 Russia; e-mail: legalov@ngs.ru

INTRODUCTION

To the family Rhynchitidae belong species, which use any other substratum for laying of eggs and species, which roll leaves into tubes for the larvae. The majority of the species belongs to the first group. Taxa are usually difficult to distinguish. The armament of the endophallus is a good character for many genera. Rhynchitidae are distributed over all natural zones. However, the centre of biodiversity is located in the Oriental area.

The author continuously studies the systematics of the leaf-rolling weevils of this family (Legalov, 2001 (2000), 2001, 2002a, 2002b, 2002c, 2003a, 2003b, 2004a, 2004b, 2005, 2006a, 2006b, 2006c, 2007, 2008, 2009; Legalov, Fremuth, 2002; Legalov, Korotyaev, 2006). The author has studied recent material (including type specimens). Results of this work are presented in this paper.

MATERIAL AND METHODS

Types and specimens are kept in the following collections and museums: BMNH – The Natural History Museum (United Kingdom: London); CKJU – P. Kresl Collection (Czech Republic: Janovice nad Uhlavou); DEI – Deutsches Entomologisches Institut (Germany: Müncheberg); ISNB – Institut Royal des Sciences Naturelles de Belgique (Belgium: Brussels); IZAS – Institute of Zoology, Academia Sinica, (China: Beijing); MCSN – Museo Civico di Storia Naturale «Giacomo Doria» (Italy: Genova); MRAC – Musee Royal de l'Afrique Centrale (Belgium: Tervuren); NHMB – Naturhistorisches Museum (Switzerland: Basel); NMPC – National Museum of Natural History (Czech Republic: Prague); RDP – R. Dunda Collection (Czech Republic: Prague); SMTD = Staatliches Museum für Tierkunde (Germany: Dresden); ZIN – Zoological Institute of Russian Academy of Sciences (Russia: St. Petersburg); ZMAN –

Zoologisch Museum, Instituut voor Taxonomische Zoologie, Universiteit van Amsterdam (Netherlands: Amsterdam); ZMHB – Museum fur Naturkunde der Humboldt-Universität (Germany: Berlin); ZMN – Zoological Museum, Institute of Animal Systematics and Ecology (Russia: Novosibirsk); ZMUC – Zoological Museum, University of Copenhagen (Denmark: Copenhagen).

RESULTS

Family Rhynchitidae Gistel, 1848
Supertribe Rhynchititae Gistel, 1848
Tribe Auletini Desbrochers des Loges,

1908

Subtribe Auletobiina Legalov, 2001
Genus *Auletobius* Desbrochers des Loges,

1869

Subgenus *Auletobius* s. str.

Auletobius (s. str.) *diversicolor* Voss,
1939 (Figs. 1, 68-69)
Auletobius diversicolor Voss, 1939b: 388

Distribution. Zaire.

Remarks. The lectotype is designated by the author – a male from the collection MRAC with labels «Paratypus», «Musée du Congo, Haut-Uele: Moto, XI.1922, L. Burgeon», «R. Dét. H. 3890», «Lectotype *Auletobius diversicolor* Voss, A. Legalov design. 2008». Paralectotype – a female from the collection MRAC with labels «Holotypus», «Musée du Congo, Haut-Uele: Moto, 1923, L. Burgeon», «R. Dét. H. 3890», «*Auletobius diversicolor* n. sp., Det. E. Voss», «Paralectotype *Auletobius diversicolor* Voss, A. Legalov design. 2008».

Key of species similar to *Auletobius kraatzi*

1. Rostrum longer. Forehead small and sparsely punctate. Cameroon..... *A. kraatzi* Voss, 1922
- . Rostrum shorter. Forehead largely and densely punctate..... 2
2. Forehead often and more weakly punctate. Pronotum completely brown. Cameroon..... *A. mengalensis* Legalov, 2007

- . Forehead more densely and largely punctate. Apex and base of pronotum yellow. Burundi.....
..... *A. urundiensis* Voss, 1939

Auletobius (s. str.) *kraatzi* Voss, 1922 (Fig.

2)

Auletobius kraatzi Voss, 1922: 31

Distribution. Cameroon.

Remarks. The holotype was studied by the author – a female from the collection DEI with labels «N Kamerun, Joh-Albrechshühe, L. Conradt 96», «Coll. Kraatz», «Voss det.», «Holotypus», «*Auletobius Kraatzi* n. sp.», «Coll. DEI Müncheberg», «*Auletobius kraatzi* Voss», «Holotype *Auletobius kraatzi* Voss, A. Legalov det. 2008».

Auletobius (s. str.) *urundiensis* Voss, 1939

(Figs. 3, 70-71)

Auletobius urundiensis Voss, 1939b: 337

Distribution. Burundi.

Remarks. The lectotype is designated by the author – a male from the collection MRAC with labels «Holotypus», «Musée du Congo, Urundi: Kanna, 26/28.I.1926, Dr. H. Schouteden», «R. Dét. E 3890», «*Auletobius urundiensis* n. sp., Det. E. Voss», «Lectotype *Auletobius urundiensis* Voss, A. Legalov design. 2008». Paralectotype – a male from the collection MRAC with labels «Paratypus», «Musée du Congo, Urundi: Kanna, 26/28.I.1926, Dr. H. Schouteden», «R. Dét. E 3890», «Paralectotype *Auletobius urundiensis* Voss, A. Legalov design. 2008».


Subgenus *Pseudometopum* Legalov, 2003

Auletobius (*Pseudometopum*)
subgranulatus Voss, 1933 (Fig. 4)
Auletobius subgranulatus Voss, 1933b: 120

Distribution. China, India, Sri Lanka, Thailand, Vietnam.

Remarks. A specimen was studied by the author – a female from the collection DEI with labels «Ceylon, Nalanda, Dr. W. Horn», «Syntypus», «*Auletobius* (*Parauletes*) *subgranulatus* m., det. E. Voss», «Coll. DEI Müncheberg», «*Auletobius subgranulatus* Voss».

Subgenus *Auletobioides* Legalov, 2007



Figs. 1-9. Auletini gen. spp.: 1 – *Auletobius diversicolor* (male, lectotype), 2 – *A. kraatzi* (female, holotype), 3 – *A. urundiensis* (male, lectotype), 4 – *A. subgranulatus* (female), 5 – *A. pallidus* (female, lectotype), 6-7 – *Pilosauletes aurichalceus* (female, holotype), 8-9 – *Pseudominurus bananensis* (male, holotype).

Auletobius (Auletobioides) pallidus
Voss, 1933, placem.n. (Fig. 5)

Auletobius pallidus Voss, 1933b: 118

Distribution. Kenya.

Remarks. The lectotype is designated by the author – a female from the collection ZMHB with labels «Tanga D. O. Afr., IV., leg. Methner, 15», «Cotype», «*Auletobius pallidus* n. sp., Det. E. Voss», «Lectotypus *Auletobius pallidus* Voss,

det. Dr. E. Haaf 1963», «Lectotypus *Auletobius pallidus* Voss, 1933, labelled by MNHUB 2008», «Lectotype *Auletobius pallidus* Voss, A. Legalov design. 2008». Paralectotype – a female from the collection ZMHB with labels «IV.15, Nairobi bei Tanga D. O. Afr., leg. Methner», «*Auletobius pallidus* m., Det. E. Voss», «Paralectotype *Auletobius pallidus* Voss, A. Legalov design. 2008».

Previously, this species has been placed wrongly in the eponymous subgenus.

Subtribe Mandelschtamiina
Legalov, 2003

Genus *Pilosauletes* Legalov, gen.n (Figs. 6-7)

Type species: *Auletobius aurichalceus* Voss, 1939

Description. Body black-brown, pronotum and elytra dark brown, femora brown. Scapus and funicle of antennae, tibiae, tarsi, basis and apex of the femora yellowish-brown. Body with dense, long, semi-erect, silvery setae. Setae concentrated on forehead, pronotum, elytra, thorax and abdomen.

Rostrum long, 5.33 times longer than wide, 1.23 times longer than pronotum, thick, straight, weakly widened to apex, with small points, looks shagreen. Antennae located near the base of the rostrum. Eyes small, weakly convex. Forehead

wide, convex, with dense small points. Temples elongated.

Antennae long, reaching the middle of the pronotum. Scapus and 1st segment of the funicle oval. 2nd-4th segments long oval, narrow. 2nd segment much longer than 1st segment. 3rd segment much shorter than 2nd segment. 3rd and 4th segments of almost equal length. 5th and 6th segments shorter than the previous segments,

wider. 7th segment almost trapezoid, short and wide. Clava narrow, almost compact, pointed. 1st and 2nd segments of almost equal length. 2nd segment wider than 1st segment. 3rd segment little shorter than the previous segments.

Pronotum weakly elongated, 1.08 times longer than width. Postnotal groove weakly but wide. Sides very weakly rounded. Disk convex, small and very densely punctate. Greatest width before middle. Scutellum wide, trapezoid.

Elytra almost rectangular, 1.33 times longer than wide. Greatest width behind middle. Humeri weakly smoothed. Striae absent. Points small and dense. Intervals flat.

Thorax small and sparsely punctate. Metepisternum narrow.

Abdomen convex. 1st and 2nd ventrites wide. 2nd ventrite wider than 1st ventrite. 3rd and 4th ventrites narrower, narrower than 1st ventrite. 5th ventrite narrow, narrower than 4th ventrite. Pygidium convex, punctate.

Legs long. Femora widened. Tibiae almost straight, weakly widened to apex. Meso- and metatibiae shorter and stronger widened in apex than protibiae. Tarsi long. 1st segment long triangular not flattened. 2nd segment wide triangular. 3rd segment bilobed. Claval segment elongated. Claws with long teeth. Length of body: 2.7 mm.

Diagnosis. This new genus is close to *Caboverdeletus* Legalov, 2007 but differs by the elytra with striae distinct in their first third, body with thin and more sparsely semi-erect hair, eyes strongly convex and tarsi elongated.

Etymology. The name of the genus is formed by adding «pilosus» («hairy») to «auletes».

***Pilosauletes aurichalceus* (Voss, 1939),
comb.n., placem.n.** (Figs. 6-7)

Auletobius aurichalceus Voss, 1939b: 42

Distribution. Rwanda.

Remarks. The holotype was studied by the author – a female from the collection MRAC with labels «Holotypus», «Musée du Congo, Ruanda: Kissenyi, 21.XII.1925, Dr. H. Schouteden», «R. Dép. D 2321», «R. Dép. B 3415», «*Auletobius aurichalceus* n. sp., Det. E. Voss», «Holotype *Auletobius aurichalceus* Voss, A. Legalov det. 2008».

Genus *Pseudominurus* Voss, 1956

***Pseudominurus bananensis* Legalov, sp.n.** (Figs. 8-9, 72)

Holotype. Male (MRAC), «Musée du Congo, Banana, 7.VIII.1920, Dr. H. Schouteden», «R. Dép. D 3890», «*Auletobius* ? *brevihirtus* m. Det. E. Voss».

Description. Male: Body yellowish-brown, tarsi, thorax, abdomen, apex of metafemora, apex of meso- and metatibiae, head and rostrum, spot near scutellum and 1st interval, scapus, 1st segment and clava brown, with dense, short, semi-erect, light setae.

Rostrum 3.5 times longer than wide, hardly shorter than pronotum, thick, very weakly curved, weakly widened to apex, sparsely punctate. Antennae located before the middle of the rostrum. Eyes large, strongly convex. Forehead wide, convex, with small points. Temples slightly narrowed to pronotum, short.

Antennae long, reaching the middle of the pronotum. Scapus and 1st segment of the funicle oval. 2nd-4th segments long oval, narrow. 2nd segment longer than 1st segment. 5th and 6th segments shorter than the previous segments. 7th segment almost trapezoid, wider than 6th segment. Clava narrow, almost compact, pointed. 1st and 2nd segments of equal length. 3rd segment little shorter than the previous segments. Pronotum 1.05 times longer than wide, with rounded sides, narrowed to base and apex. Disk convex, small and densely punctate. Greatest width in the middle. Scutellum wide, trapezoid. Elytra almost rectangular, 1.22 times longer than wide. Greatest width behind the middle. Humeri weakly smoothed. Striae weak but distinct, with large and flat points. Intervals wide, flat.

Thorax small and sparsely punctate. Metepisternum narrow.

Abdomen convex. 1st and 2nd ventrites wide. 2nd ventrite wider than 1st ventrite. 3rd and 4th ventrites narrower, narrower than 1st ventrite. 5th ventrite narrow, narrower than 4th ventrite. Pygidium convex, punctate.

Legs long. Femora widened. Tibiae almost straight, weakly widened to apex. Meso- and metatibiae shorter and stronger widened at the apex than protibiae. Tarsi long. Protarsi more

flattened and longer than meso- and metatarsi. 1st segment long triangular. 2nd segment wide triangular. 3rd segment bilobed. Claval segment elongated. Claws with long teeth. Length of body: 2.2 mm.

Distribution. Zaire.

Diagnosis. This new species is similar to *P. loidimensis* Legalov, 2007 but differs by a wider clava of the antennae, a weaker pointed apex of the aedeagus, stronger rounded sides of the pronotum.

Etymology. The name is derived from the location «Banana» – «bananensis».

***Pseudominurus brevihirtus* (Voss, 1939)**

(Fig. 10)

Auletobius brevihirtus Voss, 1939b: 339

Distribution. Zaire.

Remarks. The lectotype is designated by the author – a female from the collection MRAC with labels «Holotypus», «Musée du Congo, Haut-Uele: Abimya, VI-VII.1925, L. Burgeon», «R. Dét. C 3415», «*Auletobius brevihirtus* n. sp., Det. E. Voss», «Lectotype *Auletobius brevihirtus* Voss, A. Legalov design. 2008».

***Pseudominurus mubendensis* Legalov, 2007** (Fig. 11)

Pseudominurus mubendensis Legalov, 2007: 49

Material. Female (MRAC), «Musée du Congo, Mongbwalu (Kilo), 1938, Mme Scheitz», «R. Dét. DD 4970», «*Auletobius brevihirtus* m. Det. E. Voss».

Distribution. Uganda, Zaire.

Remarks. This species is proven the first time for the fauna of Zaire.

Subtribe Pseudauletinina Voss, 1933

Genus *Pseudauletes* Voss, 1922

Subgenus *Eopseudauletes* Legalov, 2007

***Pseudauletes (Eopseudauletes) nitens* Voss, 1930** (Figs. 12, 73-74)

Pseudauletes nitens Voss, 1930b: 61

Distribution. Brazil.

Remarks. The lectotype is designated by the author – a male from the collection NMPC with labels «Sao Paulo, Beas. Mraz lgt., Mus.

Pragense», «*Auletobius castaneus* n. sp., Det. E. Voss», «Lectotype *Auletobius castaneus* Voss, A. Legalov. design. 2008».

Subtribe Pseudomesauletinina Legalov, 2003

Genus *Gymnauletes* Legalov, 2001, placem.n. (Figs. 13-17, 75-78)

Auletobius subgenus *Gymnauletes* Legalov, 2001: 38

Type species: *Auletes glaber* Faust, 1892

Remarks. Genera *Gymnauletes* Legalov, 2001 and *Gymnauletobius* Legalov, 2007 have been placed wrongly by the author in the subtribe Pseudomesauletinina Legalov, 2003 based on studying of females. Males, now studied by the author, have an apex of the elytra with weakly sex patches. Therefore, these genera are close to genus *Hamiltoniauletes* Legalov, 2001.

***Gymnauletes bicolor* (Voss, 1922)** (Figs. 13, 75)

Auletobius bicolor Voss, 1922: 34

Distribution. Brazil.

Remarks. The holotype was studied by the author – a male from the collection ZMHB with labels «Brasil: Nirm», «34417», «*Auletobius bicolor* m., nov. spec.», «Holotypus *Auletobius bicolor* Voss, 1922, labelled by MNHUB 2008», «Holotype *Auletobius bicolor* Voss, A. Legalov det. 2008».


***Gymnauletes castaneus* (Voss, 1930)**

(Figs. 14-15, 76-77)

Auletobius castaneus Voss, 1930b: 60

Distribution. Brazil.

Remarks. The lectotype is designated by the author – a male from the collection NMPC with labels «Sao Paulo, Beas. Mraz lgt., Mus. Pragense», «*Auletobius castaneus* n. sp., Det. E. Voss», «Lectotype *Auletobius castaneus* Voss, A. Legalov. design. 2008». Paralectotype – a male with labels «Sao Paulo, Beas. Mraz lgt., Mus. Pragense», «*Auletobius castaneus* n. sp., Det. E. Voss», «Paralectotype *Auletobius castaneus* Voss, A. Legalov. design. 2008» and a female with labels «Sao Paulo, Beas. Mraz lgt., Mus. Pragense», «*Auletobius castaneus* n. sp., Det. E. Voss».



Figs. 10-15. Auletini gen. spp.: 10 – *Pseudominurus brevihirtus* (female, lectotype), 11 – *P. mubendensis* (female), 12 – *Pseudauletes nitens* (male, lectotype), 13 – *Gymnauletes bicolor* (male, holotype), 14 – *G. castaneus* (male, lectotype), 15 – *G. castaneus* (female, paralectotype).

«Paraectotype *Auletobius castaneus* Voss, A. Legalov. design. 2008».

Gymnauletes cognatus (Voss, 1930) (Figs. 16, 78)

Auletobius cognatus Voss, 1930b: 60

Distribution. Brazil.

Remarks. The lectotype is designated by the author – a male from the collection NMPC with labels «Sao Paulo, Beas. Mraz Igt., Mus. Pragense», «*Auletobius cognatus* n. sp., Det. E. Voss», «Lectotype *Auletobius cognatus* Voss, A. Legalov. design. 2008».

Gymnauletes glaber (Faust, 1892) (Fig. 17)

Auletobius glaber Faust, 1892: 43

Distribution. Brazil, Colombia, Venezuela.

Remarks. The holotype was studied by the author - a female from the collection SMTD with labels «gold small square», «Caracas, Simon», «Coll. J. Faust, Ankauf 1900», «Staatl. Museum für Tierkunde, Dresden», «Type», «*glaber* Faust», «Holotype *Auletobius glaber* Fst., A. Legalov design. 2005» (Legalov, 2007).

Genus *Gymnauletobius*
Legalov, 2007, placem.n.

Gymnauletobius Legalov, 2007: 53

type species: *Auletobius nitidus* Voss, 1922

Remarks. *Gymnauletobius* (s. str.) *kuscheli* (Voss, 1957) and *G. (s. str.) nitidus* (Voss, 1922) from Bolivia, *G. (s. str.) nudus* (Sharp, 1890) from Colombia and Panama, *G. (s. str.) peruanus* Legalov, 2007 from Peru, *G. (Amerauletes) tibialis* (Faust, 1892) from Venezuela belong to this genus.

Genus *Yunnanuletes*
Legalov, 2007

Yunnanuletes heishuensis Legalov, 2007

Yunnanuletes heishuensis Legalov, 2007: 59

Material. Female (RDP), China, S Sichuan, Daliang Shan Mts., Zhaojue vill. env., pass Xichang – Meigu vill., 12-14.VI.1998, M. Tryzna.

Distribution. China (Sichuan, Yunnan).

Remarks. This species is proven the first time for the fauna of Sichuan.

Genus *Pseudomesauletes* Legalov, 2001

Pseudomesauletes (s. str.) *contristatus*

(Voss, 1939), comb.n., placem.n. (Figs. 18, 79)

Auletobius contristatus Voss, 1939b: 45

Distribution. Rwanda, Zaire.

Remarks. The lectotype is designated by the author – a male from the collection MRAC with labels «Holotypus», «Musée du Congo, Kivu: Tshibinda, XI.1932, L. Burgeon», «R. Dét. CC 2648», «R. Dét. D 3415», «*Auletobius callosus* Voss [Hustache det.]», «*Auletobius contristatus* n. sp., Det. E. Voss», «Lectotype *Auletobius contristatus* Voss, A. Legalov design. 2008». Specimen – a male from the collection MRAC with labels «Musée du Congo, Ruanda: Kissenyi, 21.XII.1925, Dr. H. Schouteden», «R. Dét. F 3890», «*Auletobius contristatus* m., Det. E. Voss».

Previously, this species and *P. ueleanus* have been placed wrongly in genus *Alonsoiauletes* Legalov, 2003.

***Pseudomesauletes* (s. str.) *ueleanus* (Voss, 1939), comb.n., placem.n.** (Figs. 19, 80)
Auletobius ueleanus Voss, 1939b: 46

Distribution. Guinea, Zaire.

Remarks. The lectotype is designated by the author – a male from the collection NHMB with labels «Exped. Mus. G. Frey Franz. Guinea 1951, W. Afr., leg. Bechyne», «Region Kindia, Gangan, 700 m, 26.5.51», «*Auletobius ueleanus* m.», «Lectotype *Auletobius ueleanus* Voss, A. Legalov design. 2008».

Subgenus *Rubrauletes* Legalov, 2003

Key of species similar to *Pseudomesauletes consimilis*

1. Larger (3.5-4.9 mm). Funicle of antennae wider and dark. Armament of the endophallus (fig. 84). N India, Nepal. *P. poirasi* Legalov, sp.n.
- Smaller (2.2-2.6 mm). Funicle of antennae narrower and light. 2
2. Pronotum wider with more strongly rounded sides. Armament of the endophallus (fig. 81). Yunnan. *P. consimilis*
 - Pronotum narrower with more weakly rounded sides. Armament of the endophallus (fig. 83). Yunnan, Fujian. *P. kryzhanovskyi* Legalov, sp.n.

***Pseudomesauletes* (*Rubrauletes*) *consimilis* (Voss, 1930) (Figs. 20, 25, 81)**

Auletobius consimilis Voss, 1930a: 66

Distribution. China (Yunnan).

Remarks. The lectotype is designated by the author – a male from the collection ZMHB with labels «Yun-nan sen», «*Auletobius consimilis* Voss., Voss», «Syntypus *Auletobius consimilis* Voss, 1930, labelled by MNHUB 2008», «Lectotype *Auletobius consimilis* Voss, A. Legalov design. 2008».

The close new species (*P. (R.) poirasi* Legalov, sp.n.) live in India and Nepal.

***Pseudomesauletes* (*Rubrauletes*) *kryzhanovskyi* Legalov, sp.n.** (Figs. 21, 82-83)


Holotype. Male (NMPC), China, Yunnan, Jizhshan, S slope, 1600-2300 m, 23.VII.1995, Bolm.

Paratypes. Male (ZMN), China, Fujian, near Huaan, 29.V.1998; female (IZAS), China, Yunnan, Chusun – Kunming, 6.VII.1956, Kryzhanovsky.

Description. Body black-brown. Pronotum, elytra, abdomen red-brown. Scapus and funicle of antennae, coxa, basis of femora, tibiae, occasionally clava brown. Body with semi-erect, sparsely, light, short setae.

Male: Rostrum long, 5 times longer than width, 1.28-1.38 times longer than pronotum, weakly curved, widened to apex, punctate. Antennae located on the middle of the rostrum. Eyes small, strongly convex. Forehead wide, strongly convex, with small points. Temples straight lines, short.

Antennae long, reaching the first line of the pronotum. Scapus and 1st segment of funicle oval. 2nd - 5th segments long oval, narrow. 2nd segment equal to 1st segment. 3rd segment longer than 2nd segment. 4th segment shorter than 3rd segment. 5th segment oval, short. 6th segment wide oval. 7th segment almost trapezoid, sharply transversal, wider than 6th segment. Clava narrow, almost compact, pointed. 1st and 2nd segments transversal. 2nd segment hardly longer than 1st segment. 3rd segment tear-shaped, little shorter than the previous segments. Pronotum almost campaniform, length/width = 0.94-1.06, with weakly rounded sides, weakly narrowed to the basis and apex. Disk convex, small and densely punctate. Greatest width on the middle. Scutellum trapezoid.



Figs. 16-21. Auletini gen. spp.: 16 – *Gymnauletes cognatus* (male, lectotype), 17 – *G. glaber* (female, holotype), 18 – *Pseudomesauletes contristatus* (male, lectotype), 19 – *P. ueleanus* (male, lectotype), 20 – *P. consimilis* (male, lectotype), 21 – *P. kryzhanovskyi* (male, holotype).

Elytra almost rectangular, elongated, 1.41 times longer than wide. Greatest width behind the middle. Humeri weakly smoothed. Striae reduced. Points large and deep. Intervals weakly convex. Apex of the elytra with sex patches.

Thorax small and sparsely punctate. Metepisternum narrow.

Abdomen convex. 1st and 2nd ventrites wide. 2nd ventrite wider than 1st ventrite. 3rd and 4th ventrites narrower, narrower than 1st ventrite. 5th

ventrite narrow, narrower than 4th ventrite. Pygidium convex, punctate.

Legs long. Femora widened. Tibiae almost straight, weakly widened to apex. Protibiae narrow and long. Tarsi long. Protarsi hardly more flattened and longer than meso- and metatarsi. 1st segment long triangular. 2nd segment wide triangular. 3rd segment bilobed. Claval segment elongated. Claws with long teeth. Length of body: 2.2-2.5 mm.

Female: Rostrum longer, 7.0 times longer than width, 1.56 times longer than pronotum. Pronotum more elongated, 1.13 times longer than width. Elytra stronger widened to apex, 1.31 times longer than wide, without patches. Length of body: 2.6 mm.

Distribution. China (Fujian, Yunnan).

Etymology. This new species is named in honour of O.L. Kryzhanovsky.

Pseudomesauletes (Rubrauletes) poirasi

Legalov, sp.n. (Figs. 22-23, 84)

Holotype. Male (DEI), India, U. Pradesh, Mussoorie, 10 km E Mussoorie, 20 km NE Dehradun, 2000 m, 7.X.1996, Schultz & Vock.

Paratypes. Male (DEI), male (ZMN), Nepal, Himalaya, Annapura Mts., Ulleri suedl., Ghorepani, 2000 m, 16.VI.1993, Schmidt; male (RDP), N India, Uttar Pradesh, Missoorie, Kampti-Falls, 1500 m, 8.VII.1989, A. Riedel; male (CKJU), female (CKJU), N India, Uttarakhand state, 30 km N of Rishikesh, NW of Chamba, Arakot vill. env., 1800 m, 29-31.VII.2003, Z. Kejval & M. Tryzna; male (RDP), male (RDP), India, W-Arunachal Pr., Dirang vicinity, 27°21'-23' N, 92°13'-16' E, 1500-1800 m, 1-10.VI.2004, L. Dembicky; female (ISNB), «Ind Bor., Bacon», «Coll. Castelnau, Coll. Roelofs»; female (ZIN), Nepal, Kathmandu, 14.IV.1996, P. Udovichenko.

Description. Body black. Pronotum, scutellum, elytra, abdomen and claws red-brown. 1st-4th segments of funicle partially brown. Body with semi-erect, sparsely, light, short setae.

Male: Rostrum long, 6.0-7.0 times longer than wide, 1.31-1.55 times longer than pronotum, weakly curved, weakly widened to apex, sparsely punctate. Antennae located on the rostrum middle. Eyes large, strongly convex. Forehead wide, strongly convex, with small points. Temples slightly narrowed to pronotum, short.

Antennae long, reaching the middle of the pronotum. Scapus and 1st segment of funicle oval. 2nd-5th segments long oval, narrow. 2nd segment longer than 1st segment. 4th segment shorter than 3rd segment. 5th and 6th segments oval, shorter than the previous segments. 7th segment almost trapezoid, wider than 6th segment. Clava narrow, almost compact, pointed. 1st and 2nd segments transversal. 2nd segment

hardly longer than 1st segment. 3rd segment tear-shaped, little shorter than the previous segments. Pronotum almost campaniform, length/width = 0.89-1.0, with rounded sides, narrowed to the basis and apex. Disk convex, small and densely punctate. Greatest width in the middle. Scutellum trapezoid, with narrow apex.

Elytra almost rectangular, elongated, 1.36-1.40 times longer than wide. Greatest width behind the middle. Humeri weakly smoothed. Striae reduced. Points small and deep. Intervals weakly convex. Apex of the elytra with sex patches. Thorax small and sparsely punctate. Metepisternum narrow.

Abdomen convex. 1st and 2nd ventrites wide. 2nd ventrite wider than 1st ventrite. 3rd and 4th ventrites narrower, narrower than 1st ventrite. 5th ventrite narrow, narrower than 4th ventrite. Pygidium convex, punctate.

Legs long. Femora widened. Tibiae almost straight, weakly widened to apex. Meso- and metatibiae shorter and stronger widened to apex than protibiae. Tarsi long. Protarsi more flattened and longer than meso- and metatarsi. 1st segment long triangular. 2nd segment wide triangular. 3rd segment bilobed. Claval segment elongated. Claws with long teeth. Length of body: 3.5-4.9 mm.

Female: Rostrum longer, 7.8-8.0 times longer than wide, 1.45-1.78 times longer than pronotum. Pronotum length/width = 0.96-1.14. Sides of pronotum weaker rounded. Elytra 1.27-1.36 times longer than wide, more strongly widened to apex, without patches. Protibiae shorter. Length of body: 4.2-4.9 mm.

Distribution. India, Nepal.


Etymology. This new species is named in honour of A.A. Poiras.

Subgenus *Afromesauletes* Legalov, 2003

***Pseudomesauletes (Afromesauletes) viridimicans* (Voss, 1939), placem.n. (Fig. 26)**
Auletobius togoensis f. *viridimicans* Voss, 1939b: 48

Distribution. Zaire.

Remarks. The lectotype is designated by the author – a female from the collection MRAC



Figs. 22-26. *Pseudomesauletes* spp.: 22 – *P. poirasi* (male, paratype), 23 – *P. poirasi* (female, paratype), 24 – *P. kuntzeni* (male, lectotype), 25 – *P. consimilis* (male, lectotype), 26 – *P. viridimicans* (female, lectotype).

with labels «Holotypus», «Musée du Congo, Haut-Uele: Abimya, VII.1925, L. Burgeon», «R. Dét. J 3415», «R. Dét. S 1723», «R. Dét. L 2320», «*R. punctipennis* m. var. Hustache det.», «*Auletobius viridimicans* n. sp., Det. E. Voss», «Lectotype *Auletobius togoensis* f. *viridimicans* Voss, A. Legalov design. 2008».

Previously, this species has been wrongly placed in the subgenus *Metallauletes* Legalov, 2007.

Subgenus *Metallauletes* Legalov, 2007

***Pseudomesauletes* (*Metallauletes*) *kuntzeni* (Voss, 1922)** (Figs. 24, 85)
Auletobius kuntzeni Voss, 1922: 110

Distribution. Cameroon, Guinea.

Remarks. The lectotype is designated by the author – a male from the collection ZMHB with labels «Kamerun, Joh-Albrechshühe, 11.VII.-2.VIII.98, L. Conradt S.», «*Auletobius kuntzeni* m.», «Holotypus *Auletobius kuntzeni* Voss, 1922, labelled by MNHUB 2008», «Holotype *Auletobius kuntzeni* Voss, A. Legalov det. 2008».

Tribe *Eugnaptini* Voss, 1930

Subtribe *Eugnaptina* Voss, 1930

Genus *Eugnamptobius* Voss, 1922

Eugnamptobius diversus (Voss, 1948)

(Figs. 27, 86-87)

Eugnamptus diversus Voss, 1948: 156

Distribution. China (FUJ).

Remarks. The lectotype is designated by the author – a male from the collection DEI with labels «Kuatun, 2300 m, Fukien, CHINA, 7.6.1938, J. Klapperich», «Paratype *Eugnampus diversus* det. Voss», «Coll. DEI Müncheberg», «*Eugnamptus diversus* Voss», «Lectotype *Eugnamptu diversus* Voss, A. Legalov desing. 2008».

Genus *Eugnamptus* Schoenherr, 1839

Eugnamptus inclusus Voss, 1941 (Figs.

28-29)

Eugnamptus inclusus Voss, 1941b: 142

Distribution. India (E, S), Vietnam.

Remarks. The lectotype is designated by the author – a female from the collection DEI with labels «Khasia, 1898, Coll. Kraatz», «Syntypus», «*Eugnampus (Eugnamptobius) inclusus* n. sp., Det. E. Voss», «Coll. DEI Müncheberg», «*Eugnamptus inclusus* Voss», «Lectotype *Eugnamptu inclusus* Voss, A. Legalov desing. 2008».

Previously, this species has been wrongly placed in the genus *Eugnamptobius* Voss, 1922.

Genus *Cateugnamptus* Legalov, 2007

***Cateugnamptus hirsutus* (Voss, 1924), comb.n., placem.n.** (Fig. 30)
Eugnamptus hirsutus Voss, 1924: 34

Distribution. Indonesia (Java, Sumatra).

Remarks. A specimen was studied by the author – a female from the collection ZMAN with labels «G. Papandajan, Java, Drescher, 3.1915», «coll. F.C. Drescher», «*Eugnamptus hirsutus* m.», «*Eugnamptus hirsutus* Voss, 1924, ZMAN type COLE. 1640.1».

Previously this species has been wrongly placed in the genus *Neoeugnamptus* Legalov, 2003.

Genus *Eugnamptus* Schoenherr, 1839

***Eugnamptus angustatus* (Herbst, 1797)**
(Fig. 31)

Rhynchites angustatus Herbst, 1797: 140

Anthribus collaris Fabricius, 1801: 410

Distribution. North America.

Remarks. For *Anthribus collaris* the lectotype is designated by the author – a female from the collection ZMUC with labels «*collaris*», «*Anthribus collaris*, Fabricius 26», «Lectotypus *Attelabus angulatus* F., A. Legalov desig. 2007», «*Clinolabus angulatus* (Fabricius, 1787), A. Legalov det. 2008».

Tribe *Isotheini* Scudder, 1893


Subtribe *Chonostropheina* Morimoto, 1962

Genus Chonostropheus Prell, 1924

Chonostropheus tristis (Fabricius, 1794)

(Fig. 32)

Attelabus tristis Fabricius, 1794: 454



Figs. 27-32. Eugnamptini and Isotheini gen. spp.: 27 – *Eugnamptobius diversus* (male, lectotype), 28-29 – *Eugnamptus inclusus* (female, lectotype), 30 – *Cateugnamptus hirsutus* (female), 31 – *Anthribus collaris* (female, lectotype), 32 – *Attelabus tristis* (female, lectotype).

Remarks. The lectotype is designated by the author – a female without head and pronotum from the collection ZMUC with labels «*tristis*», «Lectotypus *Attelabus tristis* F., A. Legalov desig. 2007», «*Chonostropheus tristis* (Fabricius, 1794), A. Legalov det. 2008».

Subtribe Deporaina Voss, 1929

Genus *Eusproda* Sawada, 1987

Eusproda tumida (Voss, 1938), comb.n., placem.n. (Fig. 33)

Deporaus tumidus Voss, 1938a: 107

Distribution. India (Sikkim).

Remarks. The lectotype is designated by the author – a female from the collection DEI with

labels «Sikkim, Regenzeit, H. Fruhstorfer», «Holotypus», «Coll. SEI, Eberswalde», «*Deporaus (Exrhynchites) tumidus* n.sp., Det. E. Voss», «*Deporaus tumidus* Voss», «Coll. DEI Müncheberg», «Lectotype *Deporaus tumidus* Voss, A. Legalov desing. 2008», «*Eusproda tumida* (Voss, 1938), A. Legalov det. 2008».

Previously, this species has been wrongly placed in the genus *Exrhynchites* Voss, 1930.

Genus *Scolocnemus* Kirsch, 1875

Key to the *Scolocnemus scolocnemoides* species-group

1. Aedeagus wide. Apex of aedeagus pointed. Apex of metatibiae bright. Protibiae longer and stronger curved. Pronotum wider. Abdomen light. Sumatra. *Scolocnemus sumatranaensis* Legalov, 2007

-. Aedeagus narrow. Apex of aedeagus extended. Apex of metatibiae dark. Protibiae shorter and weaker curved. Pronotum narrower. Abdomen dark. Java. *Scolocnemus scolocnemoides* (Voss, 1935)

Scolocnemus scolocnemoides (Voss, 1935) (Figs. 34, 88, 90)

Deporaus scolocnemoides Voss, 1935: 112

Material. Male (NMPC), female (NMPC), «Java Centr., Dr. J. Baum»; 8 males (RDP), 4 females (RDP), East Java, Mt. Celening, 15 km SE of Lasem, 25-26.I.1998, S. Jakl.

Distribution. Indonesia (Java).

Remarks. The lectotype is designated by the author – a male from the collection DEI with labels «Noesa Kambangan, F.C. Drescher, 22-28.III.1926», «white circle», «Syntypus», «*Deporaus scolocnemoides* n. sp., Det. E. Voss», «Coll. DEI Müncheberg», «*Deporaus scolocnemoides* Voss», «Lectotype *Deporaus scolocnemoides* Voss, A. Legalov desing. 2008».

Genus *Pseudodeporaus* Voss, 1922

Subgenus *Pseudodeporaus* s. str.

Pseudodeporaus (s. str.) *kolbei* (Voss, 1938) (Figs. 35, 89, 91)

Deporaus kolbei Voss, 1938a: 108

Distribution. Indonesia (Sulawesi), Papua New Guinea.

Type depository. BMNH, ZMHB, ZMUH.

Remarks. The lectotype is designated by the author – a male from the collection ZMHB with labels «Neu-Guinea, Kaiser Wih. Land, Hatzfeldhafen, Grabowsky S.», «75934», «187», «*Rhynchites* Faust det.», «*Deporaus (Pseudodeporaus) kolbei* n. sp.», «Syntypus *Deporaus kolbei* Voss, 1938, labelled by MNHUB 2008», «Lectotype *Deporaus kolbei* Voss, A. Legalov design. 2008». Paralectotypes – 3 males from the collection ZMHB with labels «Neu-Guinea, Kaiser Wih. Land, Hatzfeldhafen, Grabowsky S.», «Syntypus *Deporaus kolbei* Voss, 1938, labelled by MNHUB 2008», «Paralectotype *Deporaus kolbei* Voss, A. Legalov design. 2008».

Genus *Caenorhinus* C.G. Thomson, 1859

Subgenus *Orientarodepus* Legalov, 2003

Caenorhinus (Orientarodepus) minimus koreanus (Voss, 1929) (Figs. 36, 92-93)


Depasophilus koreanus Voss, 1929: 28

Deporaus pauculus Voss, 1941a: 117, syn.n.

Distribution. Southern Far East of the Russia, NE, E and SE China, Korea.

Remarks. The lectotype is designated by the author – a male from the collection DEI with labels «Erzendjanzs, Manshukuo, leg. W. Alin, 23.6.1940», «Syntypus», «*Deporaus pauculus* Voss», «Coll. DEI Müncheberg». Paralectotypes: a male with labels as lectotype; a male with labels «Erzendjanzs, Manshukuo, leg. W. Alin, 23.6.1940», «Syntypus», «*Deporaus pauculus* n.sp., Det. E. Voss», «*Deporaus pauculus* Voss», «Coll. DEI Müncheberg»; male specimen with labels «Erzendjanzs, Manshukuo, leg. W. Alin, 23.6.1940», «Voss det.», «Coll. DEI Müncheberg»; 2 females with labels «Erzendjanzs, Manshukuo, leg. W. Alin, 20.6.1940», «Voss det.», «Coll. DEI Müncheberg» and female with labels «Erzendjanzs, Manshukuo, leg. W. Alin, 16.6.1940», «*Deporaus pauculus* m., Det. E. Voss», «Coll. DEI Müncheberg».

Deporaus pauculus it is identical to the continental subspecies *Caenorhinus minimus* (Kono, 1928) and can be regarded as its synonym.



Figs. 33-38. Isotheini and Rhynchitini gen. spp.: 33 – *Eusproda tumida* (female, lectotype), 34 – *Scolocnemus scolocnemoides* (male, lectotype), 35 – *Pseudodeporaus kolbei* (male, lectotype), 36 – *Deporaus pauculus* (male, lectotype), 37 – *Caenorhinus gilviventris* (male, lectotype), 38 – *Attelabus planirostris* (female, paralectotype).

Subgenus *Flavodeporaus* Legalov, 2007

***Caenorhinus* (Flavodeporaus) *gilviventris* (Voss, 1938), placem.n. (Figs. 37, 94-95)**

Deporaus gilviventris Voss, 1938a: 105

Distribution. China (FUJ), Vietnam.

Type depository. ZMHB, ZMUH.

Remarks. The lectotype is designated by the author – a male from the collection ZMHB with labels «Tonkin, Montes Mauson, April, Mai, 2-3000', H. Fruhstorfer», «*Deporaus gilviventris* n. sp., Det. E. Voss», «Syntypus *Deporaus gilviventris* Voss, 1938, labelled by MNHUB 2008», «Lectotype *Deporaus gilviventris* Voss, A. Legalov design. 2008». Paralectotype – a male from the collection ZMHB with labels «Tonkin,

Montes Mauson, April, Mai, 2-3000', H. Fruhstorfer», «Syntypus *Deporaus gilviventris* Voss, 1938, labelled by MNHUB 2008», «Paralectotype *Deporaus gilviventris* Voss, A. Legalov design. 2008».

Previously, this species has been previously placed in *Caenorhinus incertae sedis*.

Tribe Rhynchitini Gistel, 1848

Subtribe Temnocerina Legalov, 2003

Genus *Temnocerus* Thunberg, 1815

Subgenus *Temnocerus* s. str.

Temnocerus (s. str.) *nanus* (Paykull, 1792)

(Fig. 38)

Curculio nanus Paykull, 1792: 136

Attelabus planirostris Fabricius, 1801: 425

Distribution. West and Central Palaearctic.

Remarks. For *Attelabus planirostris* the lectotype is designated by the author – a male from the collection ZMUC with labels «*planirostris*» (label in entomological box), «*Lectotypus Attelabus planirostris* F., A. Legalov design. 2008», «*Temnocerus nanus* Payk., A. Legalov det. 2008». Paralectotypes: 2 females with labels «*Paralectotypus Attelabus planirostris* F., A. Legalov desig. 2008», «*Temnocerus nanus* Payk., A. Legalov det. 2008» and 2 males with labels «*Paralectotypus Attelabus planirostris* F., A. Legalov desig. 2008», «*Temnocerus coeruleus* (Fabricius, 1798), A. Legalov det. 2008» and a female with labels «*Schilsky, 1901. = Rhynchites nanus* Payk.», «*Paralectotypus Attelabus planirostris* F., A. Legalov desig. 2008», «*Temnocerus coeruleus* (Fabricius, 1798), A. Legalov det. 2008».

Two similar species (*Temnocerus nanus* č *T. coeruleus*) have been mixed in the series of type specimens.

Subtribe Perrhynchitina Legalov, 2003

Genus *Merhynchites* Sharp, 1889

Merhynchites bicolor (Fabricius, 1775)

Curculio bicolor Fabricius, 1775: 131

Distribution. North America.

Remarks. For *C. bicolor* the paralectotype was studied by the author – a female (fig. 39) from the collection ZMUC with labels «*bicolor*»,

«Paralectotype *Curculio bicolor* F., A. Legalov det. 2008», «*Attelabus variolosus* F., A. Legalov det. 2008».

This specimen (paralectotype) from type series concerns species from other family (*Attelabus variolosus* F., Attelabidae).

Genus *Japonorhynchites* Legalov, 2003

Japonorhynchites bisulcatus (Voss, 1921)

(Fig. 40)

Rhynchites bisulcatus Voss, 1921: 283

Distribution. China (Taiwan).

Remarks. The holotype was studied by the author – a female from the collection DEI with labels «*Banshoryo – Distr., Sokutsu (Formosa), H. Sauter, 1912*», «*7.VII.1912*», «*Voss det.*», «*Holotypus*», «*Rhynchites (Merhynchites) bisulcatus m.*», «*Rhynchites bisulcatus* Voss», «*Coll. DEI Müncheberg*», «*Holotype Rhynchites bisulcatus* Voss, A. Legalov det. 2008», «*Japonorhynchites bisulcatus* (Voss, 1921), A. Legalov det. 2008».

Subtribe Anisomerinina Legalov, 2003

Genus *Exochorhynchites* Voss, 1930

Exochorhynchites decumanus (Voss, 1930) (Figs. 41-42, 96)

Coenorrhinus decumanus Voss, 1930b: 62


Distribution. S-Africa.

Remarks. The lectotype is designated by the author – a male from the collection NMPC with labels «*typus*», «*Natal*», «*Mus. Pragense, Coll. Brydl*», «*Coenorrhinus decumanus* n. sp., Det. E. Voss», «*Lectotypus Coenorrhinus decumanus* Voss, A. Legalov design. 2008», «*Exochorhynchites decumanus* Voss A. Legalov det. 2008». Paralectotype – a female from the collection NMPC with labels «*typus*», «*Natal*», «*Mus. Pragense, Coll. Brydl*», «*Paralectotype Coenorrhinus decumanus* Voss, A. Legalov design. 2008», «*Exochorhynchites decumanus* Voss A. Legalov det. 2008».

Subtribe Rhynchitina Gistel, 1848

Genus *Auletomorphus* Voss, 1923

Subgenus *Auletomorphus* s. str.



Figs. 39-44. Attelabidae and Rhynchitidae gen. spp.: 39 – *Attelabus variolosus* (female, paralectotype of *Curculio bicolor*), 40 – *Japonorhynchites bisulcatus* (female, holotype), 41 – *Exochorhynchites decumanus* (female, paralectotype), 42 – *E. decumanus* (male, lectotype), 43 – *Rhynchites bicuspis* (female, lectotype), 44 – *Pararhynchites similatus* (male, lectotype).

Auletomorphus (s. str.) *cupidio* (Pascoe, 1875) (Fig. 43)

Rhynchites cupidio Pascoe, 1875: 394

Rhynchites bicuspis Voss, 1924: 42, syn.n.

Distribution. Indonesia (Sumatra), Malaysia (Perak, Pahang), S Thailand.

Remarks. The lectotype is designated by the author – a female from the collection ZMHB with

labels «W. Sumatra, 6.XII.08, Bandar Baut b., Padang Schoede S.G.», «*bicuspis* m.», «SYNTYPUS *Rhynchites bicuspis* Voss, 1934, labelled by MNHUB 2008», «Lectotype *Rhynchites bicuspis* Voss, A. Legalov design. 2008», «*Auletomorphus cupidio* (Pascoe, 1875), A. Legalov det. 2008»; paralectotype – a female from the collection ZMAN with labels «J. B.